

YOUR COLONOSCOPY APPOINTMENT

Patient's Name: _____

Tel: _____

Date: _____ Time: _____

Doctor's Name: _____

KleanLyte^{MC}

To facilitate access to endoscopic examinations, we require that you call us at least 5 days in advance if you must postpone or cancel your appointment. This will enable someone else to have their test.

- Centre multiservices de santé et services sociaux d'Argenteuil **450-562-3761** poste 72521
- Hôpital de Mont-Laurier **819-623-1234** poste 54123
- Hôpital Laurentien **819-324-4000** poste 34344 (français) et 34345 (anglais)
- Hôpital régional de Saint-Jérôme **450-432-2777** poste 22781
- Hôpital de Saint-Eustache **450-473-6811** poste 42417

What is a colonoscopy?

- ➔ A colonoscopy is a procedure in which a doctor examines the colon to detect any inflammation, polyps or colorectal cancer.

What takes place during the examination?

- ➔ A colonoscope is a flexible tube the width of a finger that is equipped with a camera. It is inserted through the anus and gradually advanced through the entire colon up to the cecum.
- ➔ To ensure good visualization of all the walls of the colon, gas or air is injected into the intestine. This can cause cramps or a bloating feeling during the test.
- ➔ To reduce the discomfort associated with the procedure, you will be given drugs intravenously to help you relax and make you drowsy. The examination takes about 20 minutes.
- ➔ A colonoscopy allows a sampling of a suspicious or inflamed area, removal of one or some polyps (abnormal growth of the intestinal inner surface), control of a bleeding, expansion of an intestinal stricture (stenose), or removing of a foreign body.

What happens after the examination?

- ➔ After the examination, you will be kept under observation until the effects of the medications have almost completely worn off (about 30 minutes).

What are the potential complications?

- ➔ Bleeding smaller than 1/100 – 200.
- ➔ Perforation smaller than 1/1,000 – 2,000 (surgery may be needed).
- ➔ Side effects may occur due to sedatives given at the time of the examination.



Your examination will require the administration of medication. Consequently, an adult must mandatorily accompany you home, otherwise your examination will be cancel. Ideally the person should stay with you for 24 hours. **NO DRIVING FOR 24 H.**

What you need to bring on the day of the examination?

- Your health insurance card and hospital card;
- **A complete list of all your medications, as provided by your pharmacist;**
- Leave any jewelry, piercing jewelry or any other valuables at home.

Instructions for medication

Take your usual medication on the day of examination, except if there is a contraindication:

- ➔ Stop iron 7 days before examination.
- ➔ You do not have to stop taking acetylsalicylic acid (Aspirin[®], Novasen[®], Entrophen[®], Asaphen[®], Rivasas^{MD}) or nonsteroidal anti-inflammatories (Advil[®], Célébrex[®], Naproxen[®]).
- ➔ If you are taking an antiplatelet agent or an anticoagulant (blood thinners), advise us, and you will receive previous instructions if your family doctor or specialist haven't already done it.
- ➔ You must tell us if you think you are pregnant.

Instructions for diabetics

Your examination should be ideally before noon.

- ➔ The day before examination: Take your usual medication for diabetes.
- ➔ The morning of examination: **Do not take** your medication for diabetes, except slow insuline **HALF the dose.** (Lantus[®], Levemir[®], NPH, Humulin[®] N).
- ➔ **It is important** to verify you glycemia and to take clear liquids allowed until **2 hours before examination.** Afterword, if you can not drink, take sugar tablets over the counter at your drugstore to raise your glycemia if needed.

Ensuring the quality of your examination

Diet



FOODS TO BE AVOIDED DURING THE 3 DAYS PRECEDING THE TEST

- X Multigrain bread, multigrain pasta, cereals and brown rice;
- X Beans and legumes (red beans, lentils, chick peas, dry peas, etc.);
- X Seeds and nuts;
- X Fruits with small seeds (strawberries, raspberries, kiwis, watermelon, blackberries, grapes, tomatoes, etc.);
- X Chips, popcorn and coconut;
- X Corn.



ON THE DAY BEFORE THE PROCEDURE, NO SOLID FOOD IS ALLOWED DRINK ONLY CLEAR LIQUIDS (no dairy products* or dairy substitutes*)

You can eat or drink as much of the following as you like throughout the day up to 2 hours before the examination:

- ✓ Water or ice;
- ✓ Clear juice without pulp (apple, white grape or lemonade);
- ✓ Jell-O® and Popsicle® (except red or purple);
- ✓ Beef or chicken consommé or broth (without noodles or vegetables and with no solid residue);
- ✓ Herbal tea, tea or coffee (without cream, milk or milk substitute);
- ✓ Clear soft drinks (Sprite®, 7-up®, Ginger-Ale®).

* Avoid all dairy products (milk or cream) and substitutes (almond milk, soya milk, Ensure, etc.)

IMPORTANT : Only for patients who are having a gastroscopy and a colonoscopy ON THE SAME DAY.

- Fast completely after taking the second complete bag of solution.
- DO NOT DRINK ANY OTHER LIQUIDS.
- If you feel or notice a drop of your sugar level, take sugar tablets available over the counter at your drugstore.

Ensure that you hydrate adequately before, during, and after taking this product.



The EVENING (at 6:00 p.m.) before the colonoscopy

STEP
1

Pour the contents of one bottle into the mixing container provided.



STEP
2

Fill the container with water to the red fill line (473 mL or 16 oz) and drink the entire amount.



STEP
3

Drink two additional containers filled to the red fill line with water over the next hour.



Watch the video on how to prepare for a colonoscopy with KleanLyte™.

The DAY of the colonoscopy

4 hours before THE PROCEDURE: repeat steps 1 to 3 using the second bottle of KleanLyte™.

STEP
1

Pour the contents of one bottle into the mixing container provided.



STEP
2

Fill the container with water to the red fill line (473 mL or 16 oz) and drink the entire amount.



STEP
3

Drink two additional containers filled to the red fill line with water over the next hour.



Do not eat anything



Drink only clear liquids you must stop drinking 2 hours before the colonoscopy. Avoid red and purple liquids, milk and alcoholic beverages.

Precautions

- Do not drink this product before first mixing with water. Drinking undiluted solution may increase the risk of nausea, vomiting, and dehydration. It is important to drink the additional recommended amount of water to prevent dehydration.
- This product may cause ulcers of the bowel or bowel problems. Do not take other laxatives while taking this product.
- Tell your healthcare professional if you have symptoms of too much fluid loss, including vomiting, nausea, bloating, dizziness, abdominal, cramping, headache, less urine than usual, or trouble drinking clear fluid.
- Tell your healthcare professional about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements. Do not take licorice or diuretics while taking this product.